

Zinc(II), copper(II) and nickel(II) dithiocarbazate complexes: structural and photophysical properties

Marcela C. Felix¹, Stephanny V. Delgado¹, Bernardo A. Iglesias², Victor M. Deflon¹

¹Instituto de Química de São Carlos, Universidade de São Paulo, São Carlos, Brasil



²Departamento de Química, Universidade Federal de Santa Maria, Santa Maria, Brasil

E-mail: marcela.felix@usp.br

Thematic Area: Coordination Chemistry

Keywords: divalent transition metals, dithiocarbazates, fluorescence.

Thiosemicarbazones and dithiocarbazates possess similar structures and can act as ligands, stabilizing transition metal complexes. In previous investigations, palladium and platinum complexes containing a pyrene-derived thiosemicarbazone ligand were presented^{1,2}. Here, new zinc(II), copper(II) and nickel(II) complexes were obtained with a new bidentate dithiocarbazate ligand containing a pyrene fluorophore group, named benzyl-(E)-2-(pyren-1-ylmethylene)hydrazine-1-carbodithioate (H-PrDTC). The ligand and the complexes [Zn(PrDTC)₂] (**1**), [Cu(PrDTC)₂] (**2**) and [Ni(PrDTC)₂] (**3**) were prepared in good yields (around 70%) and were characterized by FT-IR, UV-Vis, and NMR (¹H and ¹³C, for **1** and **3**) or EPR (for **2**) spectroscopies, elemental analysis, and single crystal X-ray diffraction (for **1** and **3**). The X-ray diffraction analyses show that complex **1** presents a distorted tetrahedral coordination geometry while complex **3** has a square-planar structure, as depicted below, left and right, respectively. The study of the photophysical properties of the compounds showed that all present steady-state fluorescence emission in the blue region, originating from the ligand due to the presence of the pyrene chromophore group in their structure.



Acknowledgments: FAPESP, CAPES, CNPq.

References

- [1] OLIVEIRA, C. G. *et. al.* Novel tetranuclear Pd^{II} and Pt^{II} anticancer complexes derived from pyrene thiosemicarbazones. *Dalton Transactions*, v. 49, n. 28, p. 9595-9604, 2020.
- [2] OLIVEIRA, C. G. *et. al.* Palladium(II) complexes with thiosemicarbazones derived from pyrene as topoisomerase IB inhibitors. *Dalton Transactions*, v. 48, n. 44, p. 16509-16517, 2019.